

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag Tag   |
| <b>Target</b>                           | CTSR3  |
| <b>Synonyms</b>                         | CACRC  |
| <b>Description</b>                      | Human CTSR3 full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | 6~8weeks   |
| <b>Uniprot ID</b>                       | Q86XQ3   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Protein Families</b>                 | Ion Channels: Other  |
| <b>Protein Pathways</b>                 | N/A  |
| <b>Molecular Weight</b>                 | The human full length CTSR3 protein has a MW of 46.4kDa  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | Voltage-gated calcium channel that plays a central role in calcium-dependent physiological responses essential for successful fertilization, such as sperm hyperactivation, acrosome reaction and chemotaxis towards the oocyte.[UniProtKB/Swiss-Prot Function]  |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |

