Human CXCL10 Protein, hFc Tag Cat. No. PME101445



PRODUCT INFORMATION

Target	CXCL10
Synonyms	C7; IFI10; INP10; IP-10; crg-2; mob-1; SCYB10; gIP-10
Description	Recombinant human CXCL10 Protein with N- terminal human Fc tag
Delivery	In Stock
Uniprot ID	P02778
Expression Host	HEK293
Тад	N-Human Fc tag
Molecular Characterization	hFc(Glu99-Ala330) CXCL10(Val22-Pro98)
Molecular Weight	54.6 KDa alter removal of the signal peptide.
Purity	The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shippin	ng Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This antimicrobial gene encodes a chemokine of the CXC subfamily and ligand for the receptor CXCR3. Binding of this protein to CXCR3 results in pleiotropic effects, including stimulation of monocytes, natural killer and T-cell migration, and modulation of adhesion molecule expression. This gene may also be a key regulator of the 'cytokine storm' immune response to SARS-CoV-2 infection. [provided by RefSeq, Sep 2020]
Usage	Research use only
Conjugate	Unconjugated

Email: info@dimabio.com Website: www.dimabio.com



Human CXCL10 Protein, hFc Tag Cat. No. PME101445





Figure 1. Human CXCL10 Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

