

PRODUCT INFORMATION

Target	CXCL7
Synonyms	Platelet Basic Protein;PBP;C-X-C Motif Chemokine 7;Leukocyte-Derived Growth Factor;LDGF;Macrophage-Derived Growth Factor;MDGF;Small-Inducible Cytokine B7;PPBP;CTAP3;CXCL7;SCYB7;TGB1;THBGB1
Description	Recombinant Human C-X-C Motif Chemokine 7 is produced by our Mammalian expression system and the target gene encoding Ser35-Asp128 is expressed with a 6His tag at the C-terminus.
Delivery	In Stock
Uniprot ID	P02775
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	Not available
Molecular Weight	11.3 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 μm filtered solution of 20mM HAc-Nac, 150mM NaCl, pH 4.0.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Human Chemokine (C-X-C motif) Ligand 7 (CXCL7), also known as neutrophil activating peptide 2 (NAP-2), is a member of the CXC chemokines containing an ELR domain (Glu-Leu-Arg tripeptide motif). Similar to other ELR domain containing CXC chemokines, such as IL-8 and the GRO proteins, CXCL7 binds CXCR2, chemoattracts and activates neutrophils. CXCL7, Connective Tissue Activating Protein III (CTAPIII) and βthromboglobulin (βTG), are proteolytically processed carboxylterminal fragments of platelet basic protein (PBP) which is found in the alphagranules of human platelets. Although CTAPIII, βTG, and PBP represent amino-terminal extended variants of NAP2 and possess the same CXC chemokine domains, these proteins do not exhibit CXCL7/NAP2 activity. CXCL7 induces cell migration through the G-protein-linked receptor CXCR-2.
Usage	Research use only






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

