

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CXCR5
Synonyms	BLR1; CD185; MDR15
Description	Human CXCR5 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P32302
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
Molecular Weight	The human full length CXCR5 protein has a MW of 42.0 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	
Background	A multi-pass membrane protein that belongs to the CXC chemokine receptor family. It is expressed in mature B-cells and Burkitt's lymphoma. This cytokine receptor binds to B-lymphocyte chemoattractant (BLC), and is involved in B-cell migration into B-cell follicles of spleen and Peyer patches.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate CXCR5-Nanodisc 0.2 μ g Human CXCR5-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CXCR5-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CXCR5-Nanodisc is 4.190ng/ml.




Figure2. Human CXCR5-Nanodisc, Flag Tag on SDS-PAGE

