

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	ENTPD2
<b>Synonyms</b>	CD39L1; NTPDase-2
<b>Description</b>	Human ENTPD2 full length protein-synthetic nanodisc
<b>Delivery</b>	3-4 weeks
<b>Uniprot ID</b>	Q9Y5L3
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
<b>Protein Pathways</b>	Purine metabolism
<b>Molecular Weight</b>	The human full length ENTPD2 protein has a MW of 53.7 kDa
<b>Formulation &amp; Reconstitution</b>	
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is the type 2 enzyme of the ecto-nucleoside triphosphate diphosphohydrolase family (E-NTPDase). E-NTPDases are a family of ecto-nucleosidases that hydrolyze 5'-triphosphates. This ecto-ATPase is an integral membrane protein. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate ENTPD2-Nanodisc 0.2 $\mu$ g Human ENTPD2-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag ENTPD2-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ENTPD2-Nanodisc is 6.083ng/ml.



Figure2. Human ENTPD2-Nanodisc, Flag Tag on SDS-PAGE

