

PRODUCT INFORMATION

| | |
|---|---|
| Target | EPHB2 |
| Synonyms | BDPLT22;CAPB;DRT;EK5;EPHT3;ERK;Hek5;PCBC;Tyro5 |
| Description | Recombinant human EPHB2 protein with C-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | P29323 |
| Expression Host | HEK293 |
| Tag | C-6×His Tag |
| Molecular Characterization | EPHB2(Val19-Leu543) 6×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 58.9 kDa after removal of the signal peptide. The apparent molecular mass of EPHB2-His is approximately 55-70 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the Eph receptor family of receptor tyrosine kinase transmembrane glycoproteins. These receptors are composed of an N-terminal glycosylated ligand-binding domain, a transmembrane region and an intracellular kinase domain. They bind ligands called ephrins and are involved in diverse cellular processes including motility, division, and differentiation. A distinguishing characteristic of Eph-ephrin signaling is that both receptors and ligands are competent to transduce a signaling cascade, resulting in bidirectional signaling. This protein belongs to a subgroup of the Eph receptors called EphB. Proteins of this subgroup are distinguished from other members of the family by sequence homology and preferential binding affinity for membrane-bound ephrin-B ligands. Allelic variants are associated with prostate and brain cancer susceptibility. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2015] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human EPHB2 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

