

PRODUCT INFORMATION

Target	EPHB3
Synonyms	EK2; ETK2; HEK2; TYRO6
Description	Recombinant human EPHB3 Protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	P54753
Expression Host	HEK293
Tag	C-6×His tag
Molecular Characterization	EPHB3(Leu38-Pro558) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 57.8 kDa after removal of the signal peptide. The apparent molecular mass of EPHB3-His is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into two groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. This gene encodes a receptor for ephrin-B family members. [provided by RefSeq, Mar 2010]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human EPHB3 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

