

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	FFAR4
Synonyms	BMIQ10; GPR120; GPR129; GT01; O3FAR1; PGR4
Description	Human FFAR4-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q5NUL3
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length FFAR4-Strep protein has a MW of 42.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Lyophilized from PBS. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a G protein-coupled receptor (GPR) which belongs to the rhodopsin family of GPRs. The encoded protein functions as a receptor for free fatty acids, including omega-3, and participates in suppressing anti-inflammatory responses and insulin sensitizing.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate FFAR4-Strep-Nanodisc 0.2 μ g Human FFAR4-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag FFAR4-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FFAR4-Strep-nanodisc is 7.707ng/ml.

kDa M R




Figure 2. Human FFAR4-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

