

PRODUCT INFORMATION

Target	FGF2
Synonyms	BFGF; FGFB; FGF-2; HBGF-2
Description	Recombinant human FGF2 Protein with C-terminal 3×Flag tag
Delivery	In Stock
Uniprot ID	P09038
Expression Host	HEK293
Tag	C-3×Flag Tag
Molecular Characterization	FGF2(Met1-Ser288) 3×Flag tag
Molecular Weight	The protein has a predicted molecular mass of 33.7 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human FGF2 Protein, Flag Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

