

PRODUCT INFORMATION

Target	FGFb
Synonyms	Fibroblast growth factor 2;FGF-2;Basic fibroblast growth factor;bFGF;Heparin-binding growth factor 2;HBGF-2;FGF2;FGFB
Description	Recombinant Human Thermostable Fibroblast Growth Factor 2(Q65I,C96S,N111G) is produced by our E.coli expression system and the target gene encoding Met1-Ser155 is expressed.
Delivery	In Stock
Uniprot ID	BAG70264.1
Expression Host	E.coli
Tag	
Molecular Characterization	Not available
Molecular Weight	17.2 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 200mM NaCl, pH 7.5.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Fibroblast growth factors (FGF) are a family of heparin-binding secreted proteins that stimulate cell proliferation and differentiation in a wide variety of tissues. FGFs play important roles in diverse biological functions both in vivo and in vitro, including mitogenesis, cellular migration, differentiation, angiogenesis, and wound healing. Human embryonic stem cell (hESC) cultures require FGF basic (also known as FGF-2 or bFGF) in cell culture media to remain in an undifferentiated and pluripotent state. Thermostable FGF basic was engineered for enhanced stability in culture media, without modification of its biological function. FGF basic is a required component of stem cell culture media for maintaining cells in an undifferentiated state. Because FGF basic is unstable, daily media changes are needed. The thermostable FGF basic that supports a 2-day media change schedule, so no media changes are required over a weekend. This thermostable FGF basic was more stable than FGF basic in biochemical studies, and maintained cell growth, pluripotency and differentiation potential with a 2-day feeding schedule.
Usage	Research use only






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

