## PRODUCT INFORMATION

| | |
|---|---|
| **Tag** | C-Flag Tag |
| **Target** | FXYD3 |
| **Synonyms** | MAT8, PLML |
| **Description** | Human FXYD3 full length protein-synthetic nanodisc |
| **Delivery** | 6~8weeks |
| **Uniprot ID** | Q14802 |
| **Expression Host** | HEK293 |
| **Protein Families** | Ion Channels: Other |
| **Protein Pathways** | N/A |
| **Molecular Weight** | The human full length FXYD3 protein has a MW of 9.3kDa |
| **Formulation & Reconstitution** | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| **Storage & Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene belongs to a small family of FXYD-domain containing regulators of Na+/K+ ATPases which share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD, and containing 7 invariant and 6 highly conserved amino acids. This gene encodes a cell membrane protein that may regulate the function of ion-pumps and ion-channels. This gene may also play a role in tumor progression. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Oct 2008] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com