

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	FZD6
Synonyms	FZ-6, FZ6, HFZ6, NDNC1, NDNC10
Description	Human FZD6-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	O60353
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	Wnt NetPath 8,Wnt signaling,Wnt signaling and pluripotency,Cancer,Notch,Wnt Pathway,
Molecular Weight	The human full length FZD6-Strep protein has a MW of 79.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	
Background	This gene represents a member of the 'frizzled' gene family, which encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The protein encoded by this family member contains a signal peptide, a cysteine-rich domain in the N-terminal extracellular region, and seven transmembrane domains, but unlike other family members, this protein does not contain a C-terminal PDZ domain-binding motif. This protein functions as a negative regulator of the canonical Wnt/beta-catenin signaling cascade, thereby inhibiting the processes that trigger oncogenic transformation, cell proliferation, and inhibition of apoptosis. Alternative splicing results in multiple transcript variants, some of which do not encode a protein with a predicted signal peptide.[provided by RefSeq, Aug 2011]
Usage	Research use only
Conjugate	Unconjugated

