

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	FZD7
Synonyms	FzE3
Description	Human FZD7 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	O75084
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling pathway
Molecular Weight	The human full length FZD7 protein has a MW of 63.6 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD7 protein contains an N-terminal signal sequence, 10 cysteine residues typical of the cysteine-rich extracellular domain of Fz family members, 7 putative transmembrane domains, and an intracellular C-terminal tail with a PDZ domain-binding motif. FZD7 gene expression may downregulate APC function and enhance beta-catenin-mediated signals in poorly differentiated human esophageal carcinomas.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate FZD7-Nanodisc 0.2 μ g Human FZD7-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag FZD7-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FZD7-Nanodisc is 1.783ng/ml.



Figure2. Human FZD7-Nanodisc, Flag Tag on SDS-PAGE

