

**PRODUCT INFORMATION**

<b>Target</b>	GDF2
<b>Synonyms</b>	BMP9; HHT5; BMP-9
<b>Description</b>	Recombinant human GDF2 Protein with C-terminal 3×Flag tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9UK05
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-3×Flag Tag
<b>Molecular Characterization</b>	GDF2(Met1-Arg429) 3×Flag tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 50.3 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 80% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein regulates cartilage and bone development, angiogenesis and differentiation of cholinergic central nervous system neurons. Mutations in this gene are associated with hereditary hemorrhagic telangiectasia. [provided by RefSeq, Jul 2016]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human GDF2 Protein, Flag Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

