

PRODUCT INFORMATION

| | |
|---|---|
| Target | GLP2R |
| Synonyms | GLP-2-R; GLP-2R |
| Description | Human GLP2R full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | O95838 |
| Expression Host | HEK293 |
| Protein Families | Druggable Genome, GPCR, Transmembrane |
| Protein Pathways | Neuroactive ligand-receptor interaction |
| Molecular Weight | The human full length GLP2R protein has a MW of 63.0 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | A G protein-coupled receptor that is closely related to the glucagon receptor and binds to glucagon-like peptide-2 (GLP2). Signalling through GLP2 stimulates intestinal growth and increases villus height in the small intestine, concomitant with increased crypt cell proliferation and decreased enterocyte apoptosis. |
| Usage | Research use only |



ELISA assay to evaluate GLP2R-Nanodisc
0.2µg Human GLP2R-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag GLP2R-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GLP2R-Nanodisc is 2.975ng/ml.



Figure2. Human GLP2R-Nanodisc, Flag Tag on SDS-PAGE

