

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	GPR65
<b>Synonyms</b>	hTDAG8; TDAG8
<b>Description</b>	Human GPR65 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q8IYL9
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length GPR65 protein has a MW of 39.3 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Receptor for the glycosphingolipid psychosine (PSY) and several related glycosphingolipids. Plays a role in immune response by maintaining lysosome function and supporting phagocytosis-mediated intracellular bacteria clearance. May have a role in activation-induced cell death or differentiation of T-cells.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate GPR65-Nanodisc 0.2 $\mu$ g Human GPR65-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GPR65-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GPR65-Nanodisc is 7.313ng/ml.



Figure 2. WB analysis of Human GPR65-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

