

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	GPRC5C
Synonyms	RAIG-3; RAIG3
Description	Human GPRC5C-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q9NQ84
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length GPRC5C-Strep protein has a MW of 48.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a member of the type 3 G protein-coupled receptor family. Members of this superfamily are characterized by a signature 7-transmembrane domain motif. The specific function of this protein is unknown; however, this protein may mediate the cellular effects of retinoic acid on the G protein signal transduction cascade.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate GPRC5C-Strep-Nanodisc 0.2 μ g Human GPRC5C-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag GPRC5C-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GPRC5C-Strep-nanodisc is 2.291ng/ml.

kDa M R



Figure 2. Human GPRC5C-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

