Delivery



PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target GRM4

Synonyms GPRC1D, MGLUR4, mGlu4

Human GRM4-Strep full length protein-synthetic **Description**

nanodisc 6~8weeks

Uniprot ID Q14833 **Expression Host HEK293**

Storage & Shipping

Protein Families GPCR, Transmembrane, Druggable Genome,

GPCRDB Class C Metabotropic glutamate pheromone, G-Protein Coupled Receptors Signaling Pathway, **Protein Pathways**

The human full length GRM4-Strep protein has a **Molecular Weight**

MW of 101.9 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

L-glutamate is the major excitatory

neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic

neurotransmission is involved in most aspects of normal brain function and can be perturbed in

many neuropathologic conditions. The

metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction

Background mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate

phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Feb 2012]

Usage Research use only Conjugate Unconjugated





