

**PRODUCT INFORMATION**

<b>Target</b>	HCRTR1
<b>Synonyms</b>	ORXR1; OX1R; OXR1
<b>Description</b>	Human HCRTR1 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	O43613
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, ES Cell Differentiation/IPS, GPCR, Transmembrane
<b>Protein Pathways</b>	Neuroactive ligand-receptor interaction
<b>Molecular Weight</b>	The human full length HCRTR1 protein has a MW of 47.5 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein is a G-protein coupled receptor involved in the regulation of feeding behavior. The encoded protein selectively binds the hypothalamic neuropeptide orexin A. A related gene (HCRTR2) encodes a G-protein coupled receptor that binds orexin A and orexin B.
<b>Usage</b>	Research use only



### ELISA assay to evaluate HCRTR1-Nanodisc 0.2 $\mu$ g Human HCRTR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag HCRTR1-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with HCRTR1-Nanodisc is 4.139ng/ml.



Figure2. WB analysis of Human HCRTR1-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

