

PRODUCT INFORMATION

HER3 **Target**

ERBB3; FERLK; LCCS2; VSCN1; ErbB-3; c-erbB3; **Synonyms** erbB3-S; MDA-BF-1; c-erbB-3; p180-ErbB3; p45-

sErbB3; p85-sErbB3

Recombinant human HER3(544-643) Protein with **Description**

C-terminal mouse Fc tag

Delivery In Stock **Uniprot ID** P21860 **HEK293 Expression Host**

Tag C-mouse Fc tag

Molecular

Storage & Shipping

Background

HER3(Ala544-Thr643) mFc(Pro99-Lys330) Characterization

The protein has a predicted molecular mass of 36.9 kDa after removal of the signal peptide. The apparent molecular mass of HER3(544-643)-mFc **Molecular Weight**

is approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and

thawing) Lyophilized proteins are shipped at

ambient temperature.

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been

characterized. One isoform lacks the

intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul

> Email: info@dimabio.com Website: www.dimabio.com

20081

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Human HER3(544-643) Protein, mFc Tag on SDS-PAGE under reducing condition.



Email: info@dimabio.com Website: www.dimabio.com

