

PRODUCT INFORMATION

Target	ICOS
Synonyms	ICOS;CD278;AILIM;Inducible T-cell costimulator
Description	Recombinant human ICOS protein with C-terminal mouse Fc and 6×His tag
Delivery	In Stock
Uniprot ID	Q9Y6W8
Expression Host	HEK293
Tag	C-Mouse Fc and 6×His Tag
Molecular Characterization	ICOS(Glu21-Lys140) mFc(Pro99-Lys330) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 41.1 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene belongs to the CD28 and CTLA-4 cell-surface receptor family. It forms homodimers and plays an important role in cell-cell signaling, immune responses, and regulation of cell proliferation.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human ICOS Protein, mFc-His Tag on SDS-PAGE under reducing condition.

Human ICOS, mFc-His Tagged protein ELISA

0.2 μg of Human ICOS, mFc-His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 $\mu\text{g}/\text{ml}$ (100 $\mu\text{l}/\text{well}$) Human ICOS, mFc-His tagged protein (PME100026) can bind Anti-ICOS Neutralizing antibody in a linear range of 0.24-31.25ng/ml.

