

PRODUCT INFORMATION

Target	IL-8
Synonyms	Interleukin-8;IL-8;C-X-C Motif Chemokine 8;CXCL8;Emoctakin;Granulocyte Chemotactic Protein 1;GCP-1;Monocyte-Derived Neutrophil Chemotactic Factor;MDNCF;Monocyte-Derived Neutrophil-Activating Peptide;MONAP;Neutrophil-Activating Protein 1;NAP-1
Description	Recombinant Human C-X-C Motif Chemokine 8/Interleukin 8 is produced by our Mammalian expression system and the target gene encoding Glu21-Ser99 is expressed with a 6His tag at the C-terminus.
Delivery	In Stock
Uniprot ID	P10145
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	Not available
Molecular Weight	10.1 KDa
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Formulation & Reconstitution	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC family of chemokines. It is one of the initial cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a variety of disease states.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

