

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	KCMB1
Synonyms	BKbeta1, K(VCA)beta, SLO-BETA, hbeta1, hslow-beta, k(VCA)beta-1, slo-beta-1
Description	Human KCMB1 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q16558
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length KCMB1 protein has a MW of 21.8kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	MaxiK channels are large conductance, voltage and calcium-sensitive potassium channels which are fundamental to the control of smooth muscle tone and neuronal excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit and the product of this gene, the modulatory beta subunit. Intracellular calcium regulates the physical association between the alpha and beta subunits. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

