Delivery

Formulation & Reconstitution

Background



PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target MC3R

Synonyms BMIQ9, MC3, MC3-R, OB20, OQTL

DescriptionHuman MC3R-Strep full length protein-synthetic

nanodisc 6~8weeks

Uniprot ID P41968 Expression Host HEK293

Protein Families GPCR,Transmembrane,Druggable Genome,

Protein Pathways

GPCRDB Class A Rhodopsin-like, Peptide

GPCRd Metabolic and Obscitu

GPCRs, Metabolic and Obesity,

Molecular Weight

The human full length MC3R-Strep protein has a MW of 36 kDa

MW of 36 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

This gene encodes a G-protein-coupled receptor for melanocyte-stimulating hormone and adrenocyticatronic hormone that is expressed in

adrenocorticotropic hormone that is expressed in tissues other than the adrenal cortex and melanocytes. This gene maps to the same region as the locus for benign neonatal epilepsy. Mice

deficient for this gene have increased fat mass despite decreased food intake, suggesting a role for this gene product in the regulation of energy homeostasis. Mutations in this gene are

> Email: info@dimabio.com Website: www.dimabio.com

associated with a susceptibility to obesity in humans. [provided by RefSeq, Jul 2008]

Usage Research use only

Conjugate Unconjugated



