

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	MCLN1
Synonyms	MG-2, ML1, ML4, MLIV, MST080, MSTP080, TRP-ML1, TRPM-L1, TRPML1
Description	Human MCLN1-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9GZU1
Expression Host	HEK293
Protein Families	Ion Channels: Transient receptor potential
Protein Pathways	N/A
Molecular Weight	The human full length MCLN1-Strep protein has a MW of 65 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the transient receptor potential (TRP) cation channel gene family. The transmembrane protein localizes to intracellular vesicular membranes including lysosomes, and functions in the late endocytic pathway and in the regulation of lysosomal exocytosis. The channel is permeable to Ca ²⁺ , Fe ²⁺ , Na ⁺ , K ⁺ , and H ⁺ , and is modulated by changes in Ca ²⁺ concentration. Mutations in this gene result in mucopolipidosis type IV. [provided by RefSeq, Oct 2009]
Usage	Research use only
Conjugate	Unconjugated

