

PRODUCT INFORMATION

Target	MERTK
Synonyms	Tyrosine-protein kinase Mer;Proto-oncogene c-Mer;Receptor tyrosine kinase MerTK
Description	Recombinant human MERTK protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q12866
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	MERTK(Ala21-Ile505) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 78.8 kDa after removal of the signal peptide. The apparent molecular mass of MERTK-hFc is approximately 100-130 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in this gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP). [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human MERTK Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

