

## **PRODUCT INFORMATION**

MMP9 **Target** 

**Synonyms** GELB;CLG4B

Recombinant human MMP9 Protein with C-Description

terminal 6×His tag

**Delivery** In Stock **Uniprot ID** P14780 **Expression Host HEK293** Tag C-6×His Tag

Molecular

**Background** 

MMP9(Ala20-Pro469) 6×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 

48.5 kDa after removal of the signal peptide. The apparent molecular mass of MMP9-His is

approximately 55-70 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & Reconstitution

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development

reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded

by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-

> Email: info@dimabio.com Website: www.dimabio.com

associated tissue remodeling. [provided by RefSeq, Jul 2008]

Usage Research use only

Conjugate Unconjugated








Figure 1. Human MMP9, His Tag on SDS-PAGE under reducing condition.



