

PRODUCT INFORMATION

Target	MST1R
Synonyms	CD136;CDw136;NPCA3;PTK8;RON
Description	Recombinant Human MST1R Protein with C-terminal Human Fc tag
Delivery	In Stock
Uniprot ID	Q04912
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	MST1R(Glu25-Thr957) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 125.9 kDa after removal of the signal peptide. The apparent molecular mass of MST1R-hFc is approximately 130-250 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a cell surface receptor for macrophage-stimulating protein (MSP) with tyrosine kinase activity. The mature form of this protein is a heterodimer of disulfide-linked alpha and beta subunits, generated by proteolytic cleavage of a single-chain precursor. The beta subunit undergoes tyrosine phosphorylation upon stimulation by MSP. This protein is expressed on the ciliated epithelia of the mucociliary transport apparatus of the lung, and together with MSP, thought to be involved in host defense. Alternative splicing generates multiple transcript variants encoding different isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Jan 2016]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human MST1R Protein, hFc Tag on SDS-PAGE under reducing condition.

Human MST1R, hFc Tagged protein ELISA

0.2 μg of Human MST1R, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 $\mu\text{g}/\text{mL}$ (100 $\mu\text{L}/\text{well}$) Human MST1R Protein, hFc Tag (PME100821) can bind Anti-MST1R Neutralizing antibody BME100118 in a linear range of 0.73–46.88 ng/mL.

