Formulation & Reconstitution



## **PRODUCT INFORMATION**

C-Flag Tag Tag MTR1A **Target** 

**Synonyms** MEL-1A-R, MT1

Human MTR1A full length protein-synthetic **Description** 

nanodisc **Delivery** 6~8weeks **Uniprot ID** P48039 **Expression Host HEK293** 

**Protein Families** GPCR, Transmembrane, Druggable Genome,

GPCRDB Class A Rhodopsin-like, Small ligand **Protein Pathways** 

GPCRs, Cancer,

The human full length MTR1A protein has a MW of **Molecular Weight** 

39.4kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Store at -20°C to -80°C for 12 months in

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes one of two high affinity forms of a receptor for melatonin, the primary hormone secreted by the pineal gland. This receptor is a G-protein coupled, 7-transmembrane receptor that is responsible for melatonin effects on

mammalian circadian rhythm and reproductive alterations affected by day length. The receptor is **Background** an integral membrané protein that is readily

detectable and localized to two specific regions of the brain. The hypothalamic suprachiasmatic nucleus appears to be involved in circadian rhythm while the hypophysial pars tuberalis may be responsible for the reproductive effects of melatonin. [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Unconjugated Conjugate

