

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	NPFF1
Synonyms	GPR147, NPFF1, NPFF1R1, OT7T022
Description	Human NPFF1-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9GZQ6
Expression Host	HEK293
Protein Families	Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,
Molecular Weight	The human full length NPFF1-Strep protein has a MW of 47.8 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	Receptor for NPAF (A-18-F-amide) and NPFF (F-8-F-amide) neuropeptides, also known as morphine-modulating peptides. Can also be activated by a variety of naturally occurring or synthetic FMRF-amide like ligands. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system.[UniProtKB/Swiss-Prot Function]
Background	
Usage	Research use only
Conjugate	Unconjugated

