

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

NPY1R **Target**

Synonyms NPY1-R; NPYR

Human NPY1R-Strep full length protein-synthetic **Description**

nanodisc **Delivery** 6~8weeks **Uniprot ID** P25929

Expression Host HEK293

Formulation &

Reconstitution

Background

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

The human full length NPY1R-Strep protein has a **Molecular Weight**

MW of 44.4 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Lyophilized from PBS. Normally 5% - 8% trehalose is added as

protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in

subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping Lyophilized proteins are shipped at ambient

temperature.

The transmembrane protein mediates the function of neuropeptide Y (NPY), a neurotransmitter, and peptide YY (PYY), a gastrointestinal hormone. The encoded receptor undergoes fast agonist-induced internalization through clathrin-coated pits and is subsequently

recycled back to the cell membrane. Activation of

Email: info@dimabio.com Website: www.dimabio.com

Y1 receptors may result in mobilization of intracellular calcium and inhibition of adenylate

cyclase activity.

Usage Research use only

Conjugate Unconjugated

