

## **PRODUCT INFORMATION**

Tag C-Flag&Strep Tag

Target OR1A1
Synonyms OR17-7

**Description**Human OR1A1-Strep full length protein-synthetic

nanodisc

Delivery 6~8weeks

Uniprot ID Q9P1Q5

Expression Host HEK293

**Protein Families** Transmembrane, Druggable Genome,

Protein Pathways GPCRDB Class A Rhodopsin-like,

Molecular Weight

The human full length OR1A1-Strep protein has a

MW of 34.6 kDa

Formulation & Reconstitution

M Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and state.

Lyophilized from nanodisc solubilization buffer (20

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the

recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is

> Email: info@dimabio.com Website: www.dimabio.com

independent of other organisms. [provided by RefSeq, Jul 2008]

Usage Research use only
Conjugate Unconjugated

**Background** 

