

PRODUCT INFORMATION

TargetP2RY6SynonymsP2Y6DescriptionHuman P2RY6-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ15077Expression HostHEK293Protein FamiliesGPCR,Transmembrane,Druggable Genome, GPCRB Class A Rhodopsin-like,Nucleotide GPCRS,Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
DescriptionHuman P2RY6-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ15077Expression HostHEK293Protein FamiliesGPCR,Transmembrane,Druggable Genome, GPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & Shipping
DescriptionnanodiscDelivery6~8weeksUniprot IDQ15077Expression HostHEK293Protein FamiliesGPCR, Transmembrane, Druggable Genome, GPCRDB Class A Rhodopsin-like, Nucleotide GPCRs, Cancer,Protein PathwaysGPCRDB Class A Rhodopsin-like, Nucleotide GPCRs, Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
Uniprot IDQ15077Expression HostHEK293Protein FamiliesGPCR,Transmembrane,Druggable Genome, GPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & Shipping
Expression HostHEK293Protein FamiliesGPCR,Transmembrane,Druggable Genome, GPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer,Molecular WeightGPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer, The human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & Shipping
Protein FamiliesGPCR,Transmembrane,Druggable Genome,Protein PathwaysGPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & Shipping
Protein PathwaysGPCRDB Class A Rhodopsin-like,Nucleotide GPCRs,Cancer,Molecular WeightThe human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStorage & Shipping
Frotein PathwaysGPCRs,Cancer, The human full length P2RY6-Strep protein has a MW of 36.4 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Storage & ShippingStorage & ShippingStorage & Shipping
Molecular WeightMW of 36.4 kDaFormulation & ReconstitutionLyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
Formulation & ReconstitutionmM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis forStorage & ShippingStore at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
Storage & Shipping Ilyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient
BackgroundThe product of this gene belongs to the family of P2 receptors, which is activated by extracellular nucleotides and subdivided into P2X ligand-gated ion channels and P2Y G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor, which is a G-protein coupled receptor, is responsive to UDP, partially responsive to UTP and ADP, and not responsive to ATP. It is proposed that this receptor mediates inflammatory responses. Alternative splicing
Usage Research use only
Conjugate Unconjugated

