

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	PD2R
Synonyms	AS1, ASRT1, DP, DP1, PTGDR1
Description	Human PD2R-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q13258
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,Prostaglandin synthesis regulation,Small ligand GPCRs,G-Protein Coupled Receptors Signaling Pathway,
Molecular Weight	The human full length PD2R-Strep protein has a MW of 40.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the guanine nucleotide-binding protein (G protein)-coupled receptor (GPCR) superfamily. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein is reported to be a receptor for prostaglandin D2, which is a mediator of allergic inflammation and allergic airway inflammation in asthma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]
Usage	Research use only
Conjugate	Unconjugated

