

**PRODUCT INFORMATION**

<b>Target</b>	pro-BDNF
<b>Synonyms</b>	Brain-Derived Neurotrophic Factor;BDNF;Abrineurin
<b>Description</b>	Recombinant Human Pro-Brain-Derived Neurotrophic Factor is produced by our E.coli expression system and the target gene encoding Ala19-Arg247(R125A,R127A,R128A) is expressed.
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P23560
<b>Expression Host</b>	E.coli
<b>Tag</b>	
<b>Molecular Characterization</b>	Not available
<b>Molecular Weight</b>	25.6 KDa
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 8.0.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The precursor form of Brain-Derived Neurotrophic Factor (pro-BDNF) interacts preferentially with the pan-neurotrophin receptor p75 (p75NTR) and vps10p domain-containing receptor sortilin and induces neuronal apoptosis, whereas mature BDNF selectively binds with high affinity to the TrkB kinase receptor and promotes the survival, growth and differentiation of neurons. As proneurotrophins and mature neurotrophins elicit opposite biological effects, Pro-BDNF cleavage in the neuronal system is regulated in a specific and cell-context dependent manner. Pro-BDNF plays important role in negative regulation of neurotrophic actions in the brain.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

