

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	PROKR1
Synonyms	GPR73; GPR73a; PK-R1; PKR1; ZAQ
Description	Human PROKR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q8TCW9
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length PROKR1 protein has a MW of 44.8 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the G-protein-coupled receptor family. The encoded protein binds to prokineticins (1 and 2), leading to the activation of MAPK and STAT signaling pathways. Prokineticins are protein ligands involved in angiogenesis and inflammation. The encoded protein is expressed in peripheral tissues such as those comprising the circulatory system, lungs, reproductive system, endocrine system and the gastrointestinal system. The protein may be involved in signaling in human fetal ovary during initiation of primordial follicle formation. Sequence variants in this gene may be associated with recurrent miscarriage.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate PROKR1-Nanodisc 0.2 μ g Human PROKR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag PROKR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with PROKR1-Nanodisc is 6.323ng/ml.



Figure2. Human PROKR1-Nanodisc, Flag Tag on SDS-PAGE

