

## PRODUCT INFORMATION

<b>Target</b>	PRTN3
<b>Synonyms</b>	ACPA;AGP7;C-ANCA;CANCA;MBN;MBT;NP-4;NP4;P29;PR-3;PR3
<b>Description</b>	Recombinant Human PRTN3 Protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P24158
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	PRTN3(Ala26-Pro256) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 26.2 kDa after removal of the signal peptide. The apparent molecular mass of PRTN3-His is approximately 25-35 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Serine protease that degrades elastin, fibronectin, laminin, vitronectin, and collagen types I, III, and IV (in vitro) (PubMed:3198760, PubMed:2033050, PubMed:28240246). By cleaving and activating receptor F2RL1/PAR-2, enhances endothelial cell barrier function and thus vascular integrity during neutrophil transendothelial migration (PubMed:23202369). May play a role in neutrophil transendothelial migration, probably when associated with CD177 (PubMed:22266279).[UniProtKB/Swiss-Prot Function]
<b>Usage</b>	Research use only






Figure 1. Human PRTN3 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

