

PRODUCT INFORMATION

Target	PTGER2
Synonyms	EP2
Description	Recombinant Human PTGER2 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P43116
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	PTGER2(Met1-Glu23) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 28.7 kDa after removal of the signal peptide. The apparent molecular mass of PTGER2-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a receptor for prostaglandin E2, a metabolite of arachidonic acid which has different biologic activities in a wide range of tissues. Mutations in this gene are associated with aspirin-induced susceptibility to asthma. [provided by RefSeq, Oct 2009]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human PTGER2 Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

