

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	QRFPR
Synonyms	AQ27; GPR103; SP9155
Description	Human QRFPR full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q96P65
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length QRFPR protein has a MW of 49.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Receptor for the orexigenic neuropeptide QRFPR. The activity of this receptor is mediated by G proteins that modulate adenylate cyclase activity and intracellular calcium levels.
Usage	Research use only
Conjugate	Unconjugated



**ELISA assay to evaluate QRFPR-Nanodisc**  
0.2µg Human QRFPR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag QRFPR-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with QRFPR-Nanodisc is 5.771ng/ml.



Figure 2. Human QRFPR-Nanodisc, Flag Tag on SDS-PAGE

