

PRODUCT INFORMATION

Target	ROR1
Synonyms	NTRKR1; dJ537F10.1
Description	Recombinant human ROR1(30-164) Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q01973
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	ROR1(Gln30-Tyr164) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 41.1 kDa after removal of the signal peptide. The apparent molecular mass of ROR1(30-164)-hFc is approximately 35-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a receptor tyrosine kinase-like orphan receptor that modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jun 2012]
Usage	Research use only






Figure 1. Human ROR1(30-164) Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMA BIO CONFIDENTIAL

