

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag&Strep Tag
<b>Target</b>	SCNNA
<b>Synonyms</b>	BESC2, ENaCa, ENaCalpha, LIDLS3, SCNEA, SCNN1
<b>Description</b>	Human SCNNA-Strep full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	P37088
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Other
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length SCNNA-Strep protein has a MW of 75.7 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Nonvoltage-gated, amiloride-sensitive, sodium channels control fluid and electrolyte transport across epithelia in many organs. These channels are heteromeric complexes consisting of 3 subunits: alpha, beta, and gamma. This gene encodes the alpha subunit, and mutations in this gene have been associated with pseudohypoaldosteronism type 1 (PHA1), a rare salt wasting disease resulting from target organ unresponsiveness to mineralocorticoids. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Apr 2009]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

