

**PRODUCT INFORMATION**

<b>Target</b>	SIGLEC5
<b>Synonyms</b>	CD33L2;CD170;OB-BP2;OBBP2;SIGLEC-5
<b>Description</b>	Recombinant Human SIGLEC5 Protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	O15389
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	SIGLEC5(Glu17-Thr434) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 47.2 kDa after removal of the signal peptide. The apparent molecular mass of SIGLEC5-His is approximately 55-100 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. These cell surface lectins are characterized by structural motifs in the immunoglobulin (Ig)-like domains and sialic acid recognition sites in the first Ig V set domain. The encoded protein is a member of the CD33-related subset of Siglecs and inhibits the activation of several cell types including monocytes, macrophages and neutrophils. Binding of group B Streptococcus (GBS) to the encoded protein plays a role in GBS immune evasion. [provided by RefSeq, Feb 2012]
<b>Usage</b>	Research use only






Figure 1. Human SIGLEC5 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

