

**PRODUCT INFORMATION**

<b>Target</b>	SLC2A12
<b>Synonyms</b>	GLUT8; GLUT12
<b>Description</b>	Human SLC2A12 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q8TD20
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length SLC2A12 protein has a MW of 67.0 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	SLC2A12 belongs to a family of transporters that catalyze the uptake of sugars through facilitated diffusion. This family of transporters show conservation of 12 transmembrane helices as well as functionally significant amino acid residues.
<b>Usage</b>	Research use only



### ELISA assay to evaluate SLC2A12-Nanodisc 0.2 $\mu$ g Human SLC2A12-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag SLC2A12-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with SLC2A12-Nanodisc is 10.17ng/ml.

kDa M R



Figure2. Human SLC2A12-Nanodisc, Flag Tag on SDS-PAGE

