

PRODUCT INFORMATION

C-Flag Tag Tag STING1 **Target**

ERIS; hMITA; hSTING; MITA; MPYS; NET23; SAVI; **Synonyms**

STING; STING-beta; TMEM173

Human STING1 full length protein-synthetic Description

nanodisc

Delivery In Stock **Uniprot ID Q86WV6** HEK293 **Expression Host**

Protein Families Transmembrane

Cytosolic DNA-sensing pathway, RIG-I-like **Protein Pathways**

receptor signaling pathway

The human full length STING1 protein has a MW Molecular Weight

of 42.2 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% Formulation & 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

A five transmembrane protein that functions as a major regulator of the innate immune response to viral and bacterial infections. The encoded protein is a pattern recognition receptor that detects cytosolic nucleic acids and transmits signals that activate type I interferon responses. The encoded protein has also been shown to play a role in

apoptotic signaling by associating with type II major histocompatibility complex. Mutations in this gene are the cause of infantile-onset STINGassociated vasculopathy. Alternate splicing

> Email: info@dimabio.com Website: www.dimabio.com

results in multiple transcript variants.

Usage Research use only Conjugate Unconjugated

Background





ELISA assay to evaluate STING1-Nanodisc 0.2µg Human STING1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag STING1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with STING1-Nanodisc is 5.896ng/ml.



Figure 2. Human STING1-Nanodisc, Flag Tag on SDS-PAGE

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

