

PRODUCT INFORMATION

Target	STING1
Synonyms	ERIS; hMITA; hSTING; MITA; MPYS; NET23; SAVI; STING; STING-beta; TMEM173
Description	Human STING1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q86WV6
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway
Molecular Weight	The human full length STING1 protein has a MW of 42.2 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A five transmembrane protein that functions as a major regulator of the innate immune response to viral and bacterial infections. The encoded protein is a pattern recognition receptor that detects cytosolic nucleic acids and transmits signals that activate type I interferon responses. The encoded protein has also been shown to play a role in apoptotic signaling by associating with type II major histocompatibility complex. Mutations in this gene are the cause of infantile-onset STING-associated vasculopathy. Alternate splicing results in multiple transcript variants.
Usage	Research use only



ELISA assay to evaluate STING1-Nanodisc 0.2 μ g Human STING1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag STING1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with STING1-Nanodisc is 5.896ng/ml.



Figure2. Human STING1-Nanodisc, Flag Tag on SDS-PAGE

