

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Tag</b>                              | C-Flag Tag  |
| <b>Target</b>                           | TAAR1   |
| <b>Synonyms</b>                         | TA1, TAR1, TRAR1  |
| <b>Description</b>                      | Human TAAR1 full length protein-synthetic nanodisc  |
| <b>Delivery</b>                         | 6~8weeks  |
| <b>Uniprot ID</b>                       | Q96RJ0  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Protein Families</b>                 | GPCR,Transmembrane,Druggable Genome,  |
| <b>Protein Pathways</b>                 | N/A   |
| <b>Molecular Weight</b>                 | The human full length TAAR1 protein has a MW of 39.1kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | The protein encoded by this gene is a G-protein coupled receptor activated by trace amines. The encoded protein responds little or not at all to dopamine, serotonin, epinephrine, or histamine, but responds well to beta-phenylethylamine, p-tyramine, octopamine, and tryptamine. While primarily functioning in neurologic systems, there is evidence that this gene is involved in blood cell and immunologic functions as well. This gene is thought to be intronless. [provided by RefSeq, Nov 2015] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |

