

## **PRODUCT INFORMATION**

TLR3 **Target** TLR3 **Synonyms** 

Recombinant human TLR3 protein with C-terminal Description

mouse Fc tag

**Delivery** In Stock **Uniprot ID** 015455 **Expression Host** HFK293

Tag C-Mouse Fc Tag

Molecular

**Background** 

TLR3(Ser24-Leu704) mFc(Pro99-Lys330) Characterization

The protein has a predicted molecular mass of

103.5 kDa after removal of the signal peptide. The apparent molecular mass of TLR3-mFc is **Molecular Weight** approximately 130-180 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.


The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They

recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of

the leukocytes. It recognizes dsRNA associated with viral infection, and induces the activation of NF-kappaB and the production of type I interferons. It may thus play a role in host defense against viruses. Use of alternative polyadenylation sites to generate different length transcripts has been noted for this gene. [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated








Figure 1. Human TLR3 Protein, mFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

