

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Target</b>                           | TNFRSF10A  |
| <b>Synonyms</b>                         | APO2;CD261;DR4;TRAILR-1;TRAILR1  |
| <b>Description</b>                      | Recombinant human TNFRSF10A protein with C-terminal human Fc tag   |
| <b>Delivery</b>                         | Under development  |
| <b>Uniprot ID</b>                       | O00220   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Tag</b>                              | C-Human Fc Tag   |
| <b>Molecular Characterization</b>       | TNFRSF10A (Ala24-Asn239) hFc (Glu99-Ala330)  |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 49.28 kDa after removal of the signal peptide.   |
| <b>Purity</b>                           | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. [provided by RefSeq, Jul 2008] |
| <b>Usage</b>                            | Research use only  |

