

PRODUCT INFORMATION

TargetTRPC7SynonymsTRP7DescriptionHuman TRPC7-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ9HCX4Expression HostHEK293Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HC1, 150 mM NaC1, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized from. After reconstitution in fund lyophilized form. After reconstitution in fund lyophilized form. After reconstitution in tended for use within a month, aliguot and store
DescriptionHuman TRPC7-Strep full length protein-synthetic nanodiscDelivery6~8weeksUniprot IDQ9HCX4Expression HostHEK293Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
DescriptionnanodiscnanodiscDelivery6~8weeksUniprot IDQ9HCX4Expression HostHEK293Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Uniprot IDQ9HCX4Expression HostHEK293Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Expression HostHEK293Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCI, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Protein FamiliesIon Channels: Transient receptor potentialProtein PathwaysN/AMolecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Protein Pathways N/A Molecular Weight The human full length TRPC7-Strep protein has a MW of 99.6 kDa Formulation & Reconstitution Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Molecular WeightThe human full length TRPC7-Strep protein has a MW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Formulation & ReconstitutionMW of 99.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
Formulation & Reconstitution mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not
lyophilized form. After reconstitution, if not
at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background Thought to form a receptor-activated non-selective calcium permeant cation channel. Probably is operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. Activated by diacylglycerol (DAG) (By similarity). May also be activated by intracellular calcium store depletion.[UniProtKB/Swiss-Prot Function]
Usage Research use only
Conjugate Unconjugated

