Delivery

Background



PRODUCT INFORMATION

Tag C-Flag Tag
Target TRPM7

Synonyms ALSPDC, CHAK, CHAK1, LTRPC7, LTrpC-7, TRP-

PLIK

DescriptionHuman TRPM7 full length protein-synthetic

nanodisc 6~8weeks

Uniprot ID Q96QT4
Expression Host HEK293

Protein Families Ion Channels: Transient receptor potential

Protein Pathways N/A

Molecular Weight
The human full length TRPM7 protein has a MW of

212.7kDa

Formulation & Reconstitution Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thains).

Lyophilized proteins are shipped at ambient

temperature.

This gene belongs to the melastatin subfamily of transient receptor potential family of ion channels. The protein encoded by this gene is both an ion channel and a serine/threonine protein kinase. The kinase activity is essential for the ion channel function, which serves to increase intracellular calcium levels and to help regulate magnesium ion homeostasis. The encoded protein is involved in cytoskeletal organization,

protein is involved in cytoskeletal organization, cell adhesion, cell migration and organogenesis. Defects in this gene are a cause of amyotrophic lateral sclerosis-parkinsonism/dementia complex of Guam. The gene may also be associated with defects of cardiac function. [provided by RefSeq,

Aug 2017]

Usage Research use only
Conjugate Unconjugated

Email: info@dimabio.com Website: www.dimabio.com



